

Skorohod-reflection of Brownian Paths and BES³

BÁLINT TÓTH and BÁLINT VETŐ

Institute of Mathematics
Technical University of Budapest (BME)

February 2, 2008

Dedicated to Sándor Csörgő on the occasion of his 60th birthday.

Abstract

Let $B(t)$, $X(t)$ and $Y(t)$ be independent standard 1d Brownian motions. Define $X^+(t)$ and $Y^-(t)$ as the trajectories of the processes $X(t)$ and $Y(t)$ pushed upwards and, respectively, downwards by $B(t)$, according to Skorohod-reflection. In the recent paper [8], Jon Warren proves inter alia that $Z(t) := X^+(t) - Y^-(t)$ is a three-dimensional Bessel-process. In this note, we present an alternative, elementary proof of this fact.

1 Introduction

The study of 1d Brownian trajectories pushed up or down by Skorohod-reflection on some other Brownian trajectories (running backwards in time) was initiated in [5] and motivated in [7] by the construction of the object what is today called the Brownian Web, see [3]. It turns out that these Brownian paths, reflected on one another, have very interesting, sometimes surprising properties. For further studies of Skorohod-reflection of Brownian paths on one another see also [6], [1], [8] etc. In particular, in [8] Warren considers two interlaced families of Brownian paths with paths belonging to the second family reflected off paths belonging to the first (in Skorohod's sense) and derives a determinantal formula for the distribution of coalescing Brownian motions.

A particular case of Warren's formula is the following: fix a Brownian path and let two other Brownian paths be pushed upwards and respectively downwards by Skorohod-reflection on the trajectory of the first one. The difference of the last two will be a

three-dimensional Bessel-process. In the present note, we give an alternative, elementary proof of this fact.

1.1 Skorohod-reflection

Let $T \in (0, \infty)$ and $b, x : [0, T) \rightarrow \mathbb{R}$ be continuous functions. Assume $x(0) \geq b(0)$. The construction of the following proposition is due to Skorohod. Its proof can be found either in [4] (see Lemma 2.1 in Chapter VI) or in [5] (see Lemma 2 in Section 2.1)

Proposition 1. (1) *There exists a unique continuous function $x_{b\uparrow} : [0, T) \rightarrow \mathbb{R}$ with the following properties*

- *The function $x_{b\uparrow} - b$ is non-negative.*
- *The function $x_{b\uparrow} - x$ is non-decreasing.*
- *The function $x_{b\uparrow} - x$ increases only when $x_{b\uparrow} = b$. That is*

$$\int_0^T \mathbb{1}\{x_{b\uparrow}(t) \neq b(t)\} d(x_{b\uparrow}(t) - x(t)) = 0.$$

(2) *The function $t \mapsto x_{b\uparrow}(t)$ is given by the construction*

$$x_{b\uparrow}(t) = x(t) + \sup_{0 \leq s \leq t} (x(s) - b(s))_-.$$

(3) *The map $C([0, T)) \times C([0, T)) \ni (b(\cdot), x(\cdot)) \mapsto (b(\cdot), x_{b\uparrow}(\cdot)) \in C([0, T)) \times C([0, T))$ is continuous in supremum distance.*

We call the function $t \mapsto x_{b\uparrow}(t)$ the *upwards Skorohod-reflection* of $x(\cdot)$ on $b(\cdot)$. As it is remarked in [5], the term *Skorohod-pushup* of $x(\cdot)$ by $b(\cdot)$ would be more adequate. Skorohod-reflection on paths $b(t) = \text{const.}$ plays a fundamental role in the proper formulation and proof of Tanaka's formula, see Chapter VI of [4].

The downwards Skorohod-reflection or Skorohod-pushdown is defined for continuous functions $b, y : [0, T) \mapsto \mathbb{R}$ with $y(0) \leq b(0)$ by

$$y_{b\downarrow} := -((-y)_{(-b)\uparrow}), \quad y_{b\downarrow}(t) = y(t) - \sup_{0 \leq s \leq t} (y(s) - b(s))_+.$$

Given three continuous trajectories $b, x, y : [0, T) \rightarrow \mathbb{R}$ with $y(0) \leq b(0) \leq x(0)$, the map $C([0, T)) \times C([0, T)) \times C([0, T)) \ni (b(\cdot), x(\cdot), y(\cdot)) \mapsto (b(\cdot), x_{b\uparrow}(\cdot), y_{b\downarrow}(\cdot)) \in C([0, T)) \times C([0, T)) \times C([0, T))$ is clearly continuous in supremum distance.

1.2 The result

Let $B(t)$, $X(t)$ and $Y(t)$ be independent standard 1d Brownian motions starting from 0 and define

$$X^+(t) := X_{B\uparrow}(t), \quad \widehat{X}(t) := X^+(t) - B(t), \quad (1)$$

$$Y^-(t) := Y_{B\downarrow}(t), \quad \widehat{Y}(t) := -Y^-(t) + B(t). \quad (2)$$

We are interested in the difference process

$$Z(t) := X^+(t) - Y^-(t) = \widehat{X}(t) + \widehat{Y}(t). \quad (3)$$

It is straightforward that $2^{-1/2}\widehat{X}(t)$ and $2^{-1/2}\widehat{Y}(t)$ are both standard reflected Brownian motions. They are, of course, strongly dependent.

The following fact is a particular consequence of the main results in [8]:

Theorem. *The process $2^{-1/2}Z(t)$ is BES³, that is standard 3d Bessel-process.*

$$dZ(t) = 2\frac{1}{Z(t)}dt + \sqrt{2}dW(t), \quad Z(0) = 0.$$

In the next section, we present an elementary proof of this fact.

2 Proof

2.1 Discrete Skorohod-reflection

Define the following square lattices inbedded in $\mathbb{R} \times \mathbb{R}$:

$$\mathcal{L} := \{(t, x) \in \mathbb{Z} \times \mathbb{Z} : t + x \text{ is even}\}, \quad \mathcal{L}^* := \{(t, x) \in \mathbb{Z} \times \mathbb{Z} : t + x \text{ is odd}\}. \quad (4)$$

In both of the lattices, the points (t_1, x_1) and (t_2, x_2) are connected with an edge if and only if $|t_1 - t_2| = |x_1 - x_2| = 1$. Note that $\mathcal{L}$ and $\mathcal{L}^*$ are Whitney-duals of each other.

We define the discrete analogue of the Skorohod-reflection in $\mathcal{L}$ and $\mathcal{L}^*$. Later on, we say that the function $y : [0, T] \cap \mathbb{Z} \rightarrow \mathbb{Z}$ is a *walk* in the lattice $\mathcal{L}$ or $\mathcal{L}^*$ if the consecutive elements of the sequence $(0, y(0)), (1, y(1)), \dots, (T, y(T))$ are edges in $\mathcal{L}$ or $\mathcal{L}^*$.

Let $b : [0, T] \cap \mathbb{Z} \rightarrow \mathbb{Z}$ and $x : [0, T] \cap \mathbb{Z} \rightarrow \mathbb{Z}$ be two walks in the lattices $\mathcal{L}$ and $\mathcal{L}^*$, respectively. Assume that $x(0) \geq b(0)$. An analogue of Proposition 1 holds in this case, but the proof is even easier.

Proposition 2. (1) *There is a unique walk $x_{b\uparrow} : [0, T] \cap \mathbb{Z} \rightarrow \mathbb{Z}$ in $\mathcal{L}^*$ with the following properties:*

- *The function $x_{b\uparrow} - b$ is non-negative.*
- *The function $x_{b\uparrow} - x$ is non-decreasing.*
- *The function $x_{b\uparrow} - x$ increases only when $x_{b\uparrow} = b + 1$, i.e.*

$$\sum_{t=1}^T \mathbb{1}\{x_{b\uparrow}(t) - b(t) > 1\} [(x_{b\uparrow}(t) - x(t)) - (x_{b\uparrow}(t-1) - x(t-1))] = 0.$$

(2) The function $t \mapsto x_{b\uparrow}(t)$ can be expressed as

$$x_{b\uparrow}(t) = x(t) + \sup_{s \in [0, t] \cap \mathbb{Z}} (x(s) - b(s))_- + 1.$$

We call the function $t \mapsto x_{b\uparrow}(t)$ the *discrete upwards Skorohod-reflection* of $x(\cdot)$ on $b(\cdot)$. The discrete downwards Skorohod-reflection is defined similarly. If $y : [0, T] \cap \mathbb{Z} \rightarrow \mathbb{Z}$ is a walk in $\mathcal{L}$ and $b : [0, T] \cap \mathbb{Z} \rightarrow \mathbb{Z}$ is a walk in $\mathcal{L}^*$ with $y(0) \leq b(0)$, then

$$y_{b\downarrow} := -((-y)_{(-b)\uparrow}), \quad y_{b\downarrow}(t) = y(t) - \sup_{s \in [0, t] \cap \mathbb{Z}} (y(s) - b(s))_+ - 1.$$

In this paper, we use the same notation for the discrete Skorohod-reflection and the continuous one (defined as Skorohod-reflection), but it will be always clear from the context which is the adequate one.

2.2 Approximation of reflected Brownian motions

Let $M(t)$ be a random walk on the lattice $\mathcal{L}$ with jumps from (t, x) to $(t + 1, x + 1)$ or $(t + 1, x - 1)$ with probability $1/2 - 1/2$ and $M(0) = 0$. We define the random walks $U(t)$ and $L(t)$ on $\mathcal{L}^*$ with the same transition probabilities, which are independent of each other and of $M(t)$. The initial values are $U(0) = 1$ and $L(0) = -1$. We extend our walks for non-integral values of t linearly, so the trajectories are continuous.

Since all these three random walks have steps with mean 0 and variance 1, it follows that

$$\left(\frac{M(nt)}{\sqrt{n}}, \frac{U(nt)}{\sqrt{n}}, \frac{L(nt)}{\sqrt{n}} \right) \xRightarrow{d} (B(t), X(t), Y(t)) \quad (n \rightarrow \infty). \quad (5)$$

We established earlier that the map $(b(\cdot), x(\cdot), y(\cdot)) \mapsto (b(\cdot), x_{b\uparrow}(\cdot), y_{b\downarrow}(\cdot))$ is continuous in supremum distance. From Donsker's invariance principle (see e.g. Section 7.6 of [2]), we conclude that

$$\left(\frac{M(nt)}{\sqrt{n}}, \frac{U_{M(n)\uparrow}(nt)}{\sqrt{n}}, \frac{L_{M(n)\downarrow}(nt)}{\sqrt{n}} \right) \xRightarrow{d} (B(t), X^+(t), Y^-(t)) \quad (6)$$

in distribution as $n \rightarrow \infty$. Note that we can use the discrete Skorohod-reflection to transform U and L , because the difference is only the addition of 1, which vanishes in the limit. At this point, it suffices to show that

$$2^{-1/2} \frac{U_{M(n)\uparrow}(nt) - L_{M(n)\downarrow}(nt)}{\sqrt{n}}$$

converges to a BES³-process.

For $x, y \in \mathbb{Z}^+$, we define the stochastic matrix

$$\mathbf{P}_{xy} = \frac{y}{x} \cdot \begin{cases} \frac{1}{2} & \text{if } y = x \\ \frac{1}{4} & \text{if } |y - x| = 1 \\ 0 & \text{otherwise} \end{cases}.$$

It is well known that if X_n is a homogeneous Markov-chain with transition probabilities $(\mathbf{P}_{xy})_{x,y \in \mathbb{Z}^+}$, then its diffusive limit is BES³, i.e. for every $T > 0$ the process $\sqrt{2}(n^{-1/2}X_{nt})_{0 \leq t \leq T}$ converges to a 3d Bessel-process in the Skorohod-topology as $n \rightarrow \infty$. So the proof of our theorem relies on the following

Lemma 1. $\frac{1}{2}(U_{M\uparrow}(t) - L_{M\downarrow}(t))$ is a Markov-chain and its transition matrix is $(\mathbf{P}_{xy})_{x,y \in \mathbb{Z}^+}$, where $U_{M\uparrow}$ and $L_{M\downarrow}$ are discrete Skorohod-reflections.

2.3 Markov-property of the distance of the two reflected walks

We introduce a different notation for the triple $(M, U_{M\uparrow}, L_{M\downarrow})$, which is just a linear transformation. Let $K_n := L_{M\downarrow}(n)$ be the position of the lower reflected walk. With the definition $D_n := \frac{1}{2}(U_{M\uparrow}(n) - L_{M\downarrow}(n))$, the distance of the two reflected walks is $2D_n$. $P_n := \frac{1}{2}(M(n) - L_{M\downarrow}(n) - 1)$, which means that the position of M related to the lower walk is $2P_n + 1$. The vector (K_n, D_n, P_n) is clearly a Markov-chain.

We are only interested in the coordinate D_n , which turns out to be also Markov and to have transition matrix $(\mathbf{P}_{xy})_{xy \in \mathbb{Z}^+}$. To show this, we have to determine the conditional distribution of P_n , because in certain cases it modifies the transition rules of D_n .

Lemma 2. *The following identities hold*

$$\mathbb{P}(P_n = x \mid D_0^n) = \frac{1}{D_n} \mathbb{1}(x \in \{0, 1, \dots, D_n - 1\}), \quad (7)$$

$$\mathbb{P}(D_{n+1} = y \mid D_0^n) = \mathbf{P}_{D_n y} \quad (8)$$

where D_0^n means the sequence of variables $D_0, \dots, D_n$.

Proof. The two identities (7), respectively, (8) of the lemma are proved by a common induction on n . Since $D_0 = 1$ and $P_0 = 0$, the case $n = 0$ is trivial.

For the induction step, we have to enumerate the possible transitions of the Markov-chain (K_n, D_n, P_n) . For the sake of simplicity, we only prove for $D_n = D_{n-1} - 1$, the other cases are similar. It is easy to check that the transition $(k, d, p) \rightarrow (k+1, d-1, p)$ has probability $\frac{1}{8} \mathbb{1}(p \in \{0, 1, \dots, d-2\})$, this will be called type *A* events. Type *B* events are the transitions $(k, d, p) \rightarrow (k+1, d-1, p-1)$, which happen with probability $\frac{1}{8} \mathbb{1}(p \in \{1, 2, \dots, d-1\})$. No other cases give $d \rightarrow d-1$.

Proof of (7): Let $x, y \in \mathbb{Z}^+$. We suppose that $y = D_{n-1} - 1$.

$$\begin{aligned}
\mathbb{P}(P_n = x \mid D_n = y, D_0^n) &= \\
&= \sum_{z \in \mathbb{Z}} \mathbb{P}(P_n = x \mid P_{n-1} = z, D_n = y, D_0^{n-1}) \mathbb{P}(P_{n-1} = z \mid D_n = y, D_0^{n-1}) \\
&= \sum_{z \in \mathbb{Z}} \frac{\mathbb{P}(P_n = x, D_n = y \mid P_{n-1} = z, D_0^{n-1})}{\mathbb{P}(D_n = y \mid P_{n-1} = z, D_0^{n-1})} \mathbb{P}(P_{n-1} = z \mid D_n = y, D_0^{n-1}) \\
&= \sum_{z=x}^{x+1} \mathbb{P}(P_n = x, D_n = y \mid P_{n-1} = z, D_0^{n-1}) \frac{\mathbb{P}(P_{n-1} = z \mid D_0^{n-1})}{\mathbb{P}(D_n = y \mid D_0^{n-1})} \\
&= \mathbb{P}(P_n = x, D_n = y \mid P_{n-1} = x, D_0^{n-1}) \frac{\mathbb{P}(P_{n-1} = x \mid D_0^{n-1})}{\mathbb{P}(D_n = y \mid D_0^{n-1})} \\
&\quad + \mathbb{P}(P_n = x, D_n = y \mid P_{n-1} = x+1, D_0^{n-1}) \frac{\mathbb{P}(P_{n-1} = x+1 \mid D_0^{n-1})}{\mathbb{P}(D_n = y \mid D_0^{n-1})} \\
&= \frac{1}{8} \mathbb{1}(x \in \{0, 1, \dots, D_{n-1} - 2\}) \frac{\frac{1}{D_{n-1}} \mathbb{1}(x \in \{0, 1, \dots, D_{n-1} - 1\})}{\frac{1}{4} \frac{D_{n-1}-1}{D_{n-1}}} \\
&\quad + \frac{1}{8} \mathbb{1}(x \in \{0, 1, \dots, D_{n-1} - 2\}) \frac{\frac{1}{D_{n-1}} \mathbb{1}(x \in \{-1, 0, \dots, D_{n-1} - 2\})}{\frac{1}{4} \frac{D_{n-1}-1}{D_{n-1}}} \\
&= \frac{1}{D_{n-1} - 1} \mathbb{1}(x \in \{0, \dots, D_{n-1} - 2\}) = \frac{1}{y} \mathbb{1}(x \in \{0, 1, \dots, y - 1\}).
\end{aligned} \tag{9}$$

First, we used the law of total probability and the definition of conditional probability and the identity $\mathbb{P}(E|F)/\mathbb{P}(F|E) = \mathbb{P}(E)/\mathbb{P}(F)$ on a conditional probability space. As remarked at the beginning of this proof, there are only two cases to reduce the value of D , so the sum has only two terms. Then, we used both inductual hypotheses to evaluate the conditional probabilities. The remaining steps are obvious.

Proof of (8): We spell out the proof for $D_{n+1} = D_n - 1$, the cases $D_{n+1} = D_n$ and $D_{n+1} = D_n + 1$ are similar.

$$\begin{aligned}
\mathbb{P}(D_{n+1} = D_n - 1 \mid D_0^n) &= \\
&= \sum_{x=0}^{D_n-1} \mathbb{P}(D_{n+1} = D_n - 1 \mid P_n = x, D_0^n) \mathbb{P}(P_n = x \mid D_0^n) \\
&= \sum_{x=0}^{D_n-1} \left(\frac{1}{8} \mathbb{1}(x \in \{0, 1, \dots, D_n - 2\}) + \frac{1}{8} \mathbb{1}(x \in \{1, 2, \dots, D_n - 1\}) \right) \frac{1}{D_n} \\
&= \frac{1}{4} \frac{D_n - 1}{D_n} = \mathbf{P}_{D_n(D_n-1)}.
\end{aligned} \tag{10}$$

In the second step, only type A and B events can cause the transition $D_{n+1} = D_n - 1$. We applied the first part of this lemma to evaluate the second conditional probability factor. □

As a consequence, we see that the distribution of D_{n+1} conditioned on D_0^n depends only on D_n , which means that D_n is a Markov-chain with transition matrix $(\mathbf{P}_{xy})_{xy}$. From this, the assertion of the theorem follows.

Acknowledgement: The result was originally proved without knowledge of Jon Warren's work. We thank Wendelin Werner for drawing our attention to the existence of the paper [8]. The authors' research is partially supported by the OTKA (Hungarian National Research Fund) grants K 60708 (for B.T. and B.V.) and TS 49835 (for B.V.).

References

- [1] K. Burdzy, D. Nualart: Brownian motion reflected on Brownian motion. *Probability Theory and Related Fields* **122**: 471-493 (2002)
- [2] R. Durrett: *Probability: Theory and Examples*. Second Edition, Duxbury Press, 1995
- [3] L.R.G. Fontes, M. Isopi, C.M. Newman, K. Ravishankar: The Brownian Web: characterization and convergence. *Annals of Probability*, **32**: 2857-2883 (2004)
- [4] D. Revuz, M. Yor: *Continuous Martingales and Brownian Motion*. Third Edition, Springer, 1999
- [5] F. Soucaliuc, B. Tóth, W. Werner: Reflection and coalescence between independent one-dimensional Brownian paths. *Annales de l'Institut Henri Poincaré – Probabilités et Statistiques*, **36**: 509-545 (2000)
- [6] F. Soucaliuc, W. Werner: A note on reflecting Brownian motion. *Electron. Comm. Probab.* **7**: 117-122 (2002)
- [7] B. Tóth, W. Werner: The true self-repelling motion. *Probability Theory and Related Fields*, **111**: 375-452 (1998)
- [8] J. Warren: Dyson's Brownian motions, intertwining and interlacing. *Electronic Journal of Probability*, **12**: 573-590 (2007)

Address of authors:
 INSTITUTE OF MATHEMATICS
 TECHNICAL UNIVERSITY OF BUDAPEST
 EGRY JÓZSEF U. 1
 H-1111 BUDAPEST, HUNGARY
 e-mail:{balint,vetob}@math.bme.hu